

Talk presented at STATPHY19, China, 1995

One Dimensional Lattice Models of Electrons with r^{-2} Hopping and Exchange

Ch. Gruber and D. F. Wang

Institut de Physique Théorique

École Polytechnique Fédérale de Lausanne

PHB-Ecublens, CH-1015 Lausanne-Switzerland.

This talk reviews some recent progresses in the study of low dimensional electron systems of strong correlations.

PACS number: 71.30.+h, 05.30.-d, 74.65+n, 75.10.Jm

I. INTRODUCTION: INTEREST OF 1D SYSTEMS

The study of one dimensional systems is almost as old as statistical mechanics. In the early days it might have been considered to be just a “warm up” exercise necessary before going to the study of the real 3-dimensional world. However, in the last decades, the study of strongly correlated electron systems has been very active to understand general properties of condensed matter, such as metal-insulator transition and high temperature superconductor, and 1D systems are expected to throw some light on the mechanisms underlying such phenomena. More recently, the discovery of high temperature superconductivity, and the suggestion by Anderson that the 2D Hubbard model could be appropriate to describe these new materials, have stimulated considerable interest in the study of electron systems in low dimensions. This is due to the possibility that the normal state of these 2D superconducting materials may share some properties of the 1D interacting electron systems, the non-Fermi-liquid behaviors. Finally, we wish to mention that today one can realize 1D conductors and 1D magnets experimentally.

II. 1D LATTICE MODELS: A SHORT REVIEW

The systems usually considered consist of identical electrons with internal degree of freedom (or “spins”) on a finite lattice $\Lambda = \{x\}$, with sites x in R . The internal degrees of freedom is denoted by σ , with $\sigma = 1, 2, \dots, \nu$. The electrons interact with each other and can also interact with “classical particles” (or f-electrons in the study of rare earth materials, and mixed valence systems), as well as with “impurity spins” (e.g. Kondo models), located on the vertices of the lattice.

The electrons are described by creation and annihilation operators $a_{x\sigma}^\dagger, a_{x\sigma}$, which satisfy the usual anti-commutation relations $\{a_{x\sigma}^\dagger, a_{y\sigma'}\} = \delta_{xy}\delta_{\sigma\sigma'}, \dots$; the classical particles are described by a random variables ω_x , taking for example the values 1 and 0 to describe the presence or absence of a particle at the site x , and the impurity spin at x is denoted by $\vec{S}_x^f$.

The Hamiltonian of physical interest are typically of the form $H = T + W$, where the kinetic part T is given by

$$T = - \sum_{x,y} \sum_{\sigma,\sigma'} t_{xy}^{\sigma\sigma'} a_{x\sigma}^\dagger a_{y\sigma'} + \sum_{x,y} \sum_{\sigma,\sigma'} \tilde{t}_{xy}^{\sigma\sigma'} a_{x\sigma}^\dagger a_{y\sigma'} [n_{x,-\sigma} + n_{y,-\sigma'} - \gamma n_{x,-\sigma} n_{y,-\sigma'}], \quad (1)$$

where $t_{xy}^{\sigma\sigma}$ is the usual hopping matrix, $t_{xy}^{\sigma\sigma'}, \sigma \neq \sigma'$, takes into account spin orbit coupling, $\tilde{t}_{xy}^{\sigma\sigma'}$, takes into account bond-charge repulsion (written here for $\nu = 2$ with $\sigma = \pm 1$). The potential part W is given by

$$W = \sum_{x,y} \sum_{\sigma,\sigma'} U_{xy}^{\sigma\sigma'} n_{x\sigma} n_{y\sigma'} + \sum_{x,y} J_{x,y} \vec{S}_x \cdot \vec{S}_y + \tilde{U} \sum_x \omega_x n_x + \tilde{J} \sum_x \vec{S}_x^f \cdot \vec{S}_x, \quad (2)$$

where $U, J, \tilde{U}, \tilde{J}$ are coupling constants (or interaction potentials), $n_{x\sigma} = a_{x\sigma}^\dagger a_{x\sigma}$, and $n_x = \sum_\sigma n_{x\sigma}$.

In this context, the simplest model is the Falicov-Kimball model which describes spinless electrons ($\nu = 1$), hopping between nearest neighbours and interacting only with the classical particles, [1] i. e.:

$$H_{F.K.} = -t \sum_{\langle x,y \rangle} a_x^\dagger a_y + \tilde{U} \sum_x \omega_x n_x. \quad (3)$$

Other typical short range models are

- The Hubbard model ($\nu = 2, \sigma = \uparrow$ or $\downarrow$) [4],

$$H_H = - \sum_{\langle x,y \rangle} \sum_{\sigma} t^{\sigma\sigma} a_{x\sigma}^\dagger a_{y\sigma} + \tilde{U} \sum_x n_{x\uparrow} n_{x\downarrow}. \quad (4)$$

In the ordinary Hubbard model $t^{\uparrow\uparrow} = t^{\downarrow\downarrow} = t$. On the other hand, taking $t^{\uparrow\uparrow} = t$ and $t^{\downarrow\downarrow} = 0$, one recovers the F.K. model with $\sigma = \downarrow$, corresponding to the classical particles.

- The Hubbard model with spin flip for which $t_{xy}^{\sigma\sigma'} = t$ for nearest neighbours, zero otherwise, and $\tilde{t}_{xy}^{\sigma\sigma'} = 0$. Introducing the operators [3]

$$\begin{aligned} A_x^\dagger &= \frac{1}{\sqrt{2}} (a_{x\uparrow}^\dagger + a_{x\downarrow}^\dagger) \\ B_x^\dagger &= \frac{1}{\sqrt{2}} (a_{x\uparrow}^\dagger - a_{x\downarrow}^\dagger), \end{aligned} \quad (5)$$

one recovers again the F. K. model, with B describing the classical particles.

- The generalized Hubbard model, where

$$H = H_H + V \sum_{\langle x,y \rangle} n_x n_y + \tilde{U} \sum_x \omega_x n_x. \quad (6)$$

· The Montorsi-Rasetti model, where $\nu = 2$ and $t_{\langle x,y \rangle}^{\sigma\sigma'} = \tilde{t}_{\langle x,y \rangle}^{\sigma\sigma'} = t$ for nearest neighbors. Introducing the operators $A_x^\dagger$ and $B_x^\dagger$ as above, one obtain the Hirsch model with B immobile.

One the other hand, several long range models have been studied, e.g.,

- $H = -\sum_{x \neq y} \sum_{\sigma} t_{xy} a_{x\sigma}^\dagger a_{y\sigma} + U \sum_x n_x n_x$, with $t_{xy} = it(-1)^{(x-y)}/d_{xy}$, and $d_{xy} = \frac{L}{\pi} \sin[\frac{\pi(x-y)}{L}]$.
- “Kondo Lattice models”, $H = \sum_{x \neq y} \sum_{\sigma} t_{xy} a_{x\sigma}^\dagger a_{y\sigma} + J \sum_x \vec{S}_x^f \cdot \vec{S}_x$, with $t_{xy} = t$. [16]

In 1969, Lieb and Wu obtained the exact solution for the 1D Hubbard model using Bethe-ansatz techniques [2]. Later, motivated by the work of Calogero, Sutherland and Moser on the integrability of 1D electron in the continuum with r^{-2} interaction [15], Haldane and Shastry introduced the spin chain with $1/r^2$ interaction [5,6]. In the following, we shall concentrate on the t-J models with $1/r^2$ interaction and discuss various properties of these systems.

III. T-J MODELS

The t-J models discuss lattice systems of electrons with hard core, i.e., the wavefunctions for N_e electrons $\Psi(x_1\sigma_1, \dots, x_{N_e}\sigma_{N_e})$, $x_i \in \Lambda$, $\sigma_i \in \{1, 2, \dots, \nu\}$ must satisfy the constrain $\Psi = 0$ if $x_i = x_j$ for some pair (i, j) . The Hamiltonian is

$$H_{tJ} = P_G \left\{ - \sum_{x,y} \sum_{\sigma=1}^{\nu} t_{xy} a_{x\sigma}^\dagger a_{y\sigma} + \sum_{x,y} J_{xy} [P_{xy} - (1 - n_x)(1 - n_y)] \right\} P_G, \quad (7)$$

where P_G is the Gutzwiller projector onto those states satisfying the constraint that there is at most one electron at each site, and P_{xy} is the operator which permutes the spins of the electrons at sites x and y , and which is zero if the two sites are not both occupied.

Following the usual approach we replace the original model by a new one with two types of particles F and B satisfying the constraint that there is exactly one particle at each site. The F particles are fermions with spins, described by the operators $f_{x\sigma}^\dagger, f_{x\sigma}$, $x \in \Lambda, \sigma \in \{1, 2, \dots, \nu\}$, the B particles are spinless bosons described by $b_x^\dagger, b_x$. The constraint, expressed by $b_x^\dagger b_x + \sum_{\sigma} f_{x\sigma}^\dagger f_{x\sigma} = 1$, implies that the wavefunction for N_e fermions and Q

bosons $\phi(x_1\sigma_1, \dots, x_{N_e}; y_1, \dots, y_Q)$ is zero, if $\{x_j\} \cup \{y_l\} \neq \Lambda$. In particular, $N_e + Q = N$, the number of the lattice sites.

The two models are then isomorphic with the obvious mapping

$$\begin{aligned}\Psi(x_1\sigma_1, \dots, x_{N_e}\sigma_{N_e}) &= \phi(x_1\sigma_1, \dots, x_{N_e}\sigma_{N_e}, y_1, \dots, y_Q) \\ a_{x\sigma}^\dagger &\rightarrow X_x^{\sigma 0} = f_{x\sigma}^\dagger b_x \\ a_{x\sigma} &\rightarrow X_x^{0\sigma} = b_x^\dagger f_{x\sigma}.\end{aligned}\tag{8}$$

Furthermore, introducing the operators

$$\begin{aligned}X_x^{\sigma\sigma'} &= f_{x\sigma}^\dagger f_{x\sigma'} \\ X_x^{00} &= b_x^\dagger b_x,\end{aligned}\tag{9}$$

the Hamiltonian H_{tJ} is mapped on

$$\begin{aligned}H &= P_1 \left\{ - \sum_{x,y} \sum_{\sigma} \frac{1}{2} (t_{xy} X_x^{\sigma 0} X_y^{0\sigma} - \bar{t}_{xy} X_x^{0\sigma} X_y^{\sigma 0}) + \right. \\ &\quad \left. + \sum_{xy} J_{xy} \left(\sum_{\sigma, \sigma'} X_x^{\sigma\sigma'} X_y^{\sigma'\sigma} - X_x^{00} X_y^{00} \right) \right\} P_1,\end{aligned}\tag{10}$$

where P_1 is the projector onto those states satisfying those constraints. The t-J model is supersymmetric if

$$t_{xy} = \bar{t}_{xy}, \quad J_{xy} = \frac{1}{2} t_{xy},\tag{11}$$

and in this case,

$$H = P_1 \left\{ - \sum_{x,y} \sum_{a,b} J_{xy} X_x^{ab} X_y^{ba} \theta_b \right\} P_1,\tag{12}$$

where $a, b \in \{0, 1, \dots, \nu\}$ and $\theta_0 = +1, \theta_\sigma = -1$. The operators X_x^{ab} generate a superalgebra. This superalgebra induces superrotations which mix the F and B particles, but leaves invariant $\sum_a X_x^{aa} = 1$. Finally, the supersymmetric t-J model is invariant under these superrotations. The usual $SU(2)$, supersymmetric, short range t-J model on a uniform lattice, i.e.

$$\nu = 2, \begin{cases} t_{xy} = t, J_{xy} = J = t/2, & \text{if } |x - y| = 1, \\ t_{xy} = J_{xy} = 0, & \text{otherwise} \end{cases} \quad (13)$$

is integrable, and was solved first by Sutherland in 70's. Following the works of Haldane and Shastry on the spin chain with $1/r^2$ interaction, Kuramoto and Yokoyama introduced a $SU(2)$, long range t-J model with [7]

$$\begin{cases} J_{xy} = \frac{1}{2}t_{xy} = 1/d_{xy}^2 \\ d_{xy} = \frac{L}{\pi} \sin(\frac{\pi(x-y)}{L}). \end{cases} \quad (14)$$

They obtained the ground state wavefunction away from the half-filling. Later the asymptotic energy spectrum was derived on the assumption of asymptotic scattering matrix factorization [8], and the system was identified as a free Luttinger liquid. Generalized Jastrow wavefunctions for excitations of the system were explicitly constructed with the representations of down spins and holes by Wang, Liu and Coleman, and they constructed the full energy spectrum and thermodynamics of the system in terms of more generalized Jastrow wavefunctions [9], which will be discussed in the next section.

IV. SUPERSYMMETRIC T-J MODEL WITH $1/R^2$ HOPPING AND EXCHANGE ON THE UNIFORM LATTICE

With the mapping discussed in Section 3, the eigenvalue equation $H_{tJ}|\Psi\rangle = E|\Psi\rangle$ for the electron model is mapped onto the equation $H|\phi\rangle = E|\phi\rangle$ for the F-B model with H given by Eq.(12) and $|\phi\rangle$ can be written as

$$|\phi\rangle = \sum_{\{x\}, \{y\}} \sum_{\sigma_1, \dots, \sigma_{N_e}} \phi(x_1\sigma_1, \dots, x_{N_e}\sigma_{N_e}; y_1, \dots, y_Q) \cdot f_{x_1\sigma_1}^\dagger \dots f_{x_{N_e}\sigma_{N_e}}^\dagger b_{y_1}^\dagger \dots b_{y_Q}^\dagger |0\rangle, \quad (15)$$

where N_e is the number of fermions and Q is the number of bosons on the lattice, $\{x\} \cup \{y\} = \Lambda$. Let us introduce the notation,

$$\begin{aligned} \phi(x_1\sigma_1, \dots, x_{N_e}\sigma_{N_e}; y_1, \dots, y_Q) &= \phi(\{q\}; \{\sigma\}) \\ (q_1, \dots, q_N) &= (x_1, \dots, x_{N_e}, y_1, \dots, y_Q), \quad N_e + Q = N. \end{aligned} \quad (16)$$

The eigenvalue equation for N_e fermions and Q bosons of the F-B model takes the form

$$-\frac{1}{2}[\sum_{i \neq j} d^{-2}(q_i - q_j) M_{ij}] \phi(\{q\}; \{\sigma\}) = E \phi(\{q\}; \{\sigma\}), \quad (17)$$

where M_{ij} is the operator which exchange the positions of particles i and j :

$$\begin{aligned} (M_{ij} \phi)(\{q\}; \{\sigma\}) &= \phi(\{q'\}; \{\sigma\}) \\ (q'_1, \dots, q'_i, \dots, q'_j, \dots, q'_N) &= (q_1, \dots, q_j, \dots, q_i, \dots, q_N). \end{aligned} \quad (18)$$

At this point, we can use the exchange operator formalism first introduced by Polychronakos to exhibit a complete set of constants of motion. In fact, this method is used by Fowler and Minahan to obtain the constants of the motion of the spin chain model of Haldane and Shastry (to which the t-J model reduces at half-filling). Following the idea of Fowler and Minahan [11], we define the operators

$$\begin{cases} \pi_j = \pi_j^\dagger = \sum_{k=1(\neq j)}^N \frac{z_k}{z_j - z_k} M_{jk}, & j = 1, \dots, N \\ z_j = \exp(i \frac{2\pi}{N} q_j), \end{cases} \quad (19)$$

and $I_n = \sum_{j=1}^N \pi_j^n$, $n = 0, 1, 2, \dots, \infty$. Using the facts that all sites are occupied by exactly one particle and that the lattice is invariant under translation, it was shown [11] that

$$\begin{aligned} [I_n, I_m] &= 0, \\ [H, I_n] &= 0, \quad n, m = 0, 1, 2, \dots \end{aligned} \quad (20)$$

Using the properties of the wave function under permutation it is straightforward to write out all the constants of the motion in terms of creation and annihilation operators [12].

With the integrability of the model established, the eigenfunctions can be explicitly constructed. In the $SU(2)$ case, one starts from the fully polarized up-spin state $|P\rangle$. Let M denote the number of the down spins and Q the number of the holes, then the wavefunction can be written as [9]

$$|\phi\rangle = \sum_{x,y} \phi(x,y) \prod_{\alpha} S_{x\alpha}^- \prod_i h_{yi}^+ |P\rangle, \quad (21)$$

where $\phi(x, y)$ is symmetric in $x = (x_1, \dots, x_M)$, the positions of the down-spins, and antisymmetric in $y = (y_1, \dots, y_Q)$, the positions of the holes; $S_x^- = a_{x\uparrow}^\dagger a_{x\downarrow}$, $h_y^\dagger = a_{y\uparrow}$, $S_z = (N - Q)/2 - M$. A large class of Jastrow product eigenstates of uniform motion and spin polarization are given by [9]

$$\begin{aligned}\phi(x, y, J_s, J_h) &= e^{\frac{2\pi i}{N}(J_s \sum_\alpha x_\alpha + J_h \sum_i y_i)} \phi_0(x, y) \\ \phi_0(x, y) &= \prod_{\alpha < \beta} d^2(x_\alpha - x_\beta) \prod_{i < j} d(y_i - y_j) \prod_{\alpha, i} d(x_\alpha - y_i),\end{aligned}\tag{22}$$

where the quantum numbers J_s and J_h satisfy the following constraints $|J_s - N/2| \leq N/2 - (M - 1 + Q/2)$, $|J_h - N/2| \leq N/2 - (M + Q - 1)/2$ and $|J_h - J_s| \leq (M + 1)/2$. Further, the full energy spectrum and thermodynamics can also be studied by looking at more generalized Jastrow wavefunctions [9].

V. SUPERSYMMETRIC T-J MODEL WITH R^{-2} HOPPING AND EXCHANGE ON A NON-UNIFORM LATTICE

In this section, we discuss the supersymmetric t-J model defined on a non-uniform lattice [13,14]. The sites of the lattice are given by the roots of the Hermit polynomial $H_N(x)$. It is well known that the Hermit polynomial $H_N(x)$ has N roots, all of which are real and distinct. Therefore the lattice is well defined. In this case, $J_{xy} = t_{xy}/2 = 1/(x - y)^2$. At half-filling, this t-J model reduces to the spin chain introduced by Polychronakos [10]. With the super-algebra presentation, and with the permutation symmetry of the wavefunction, the t-J model Hamiltonian can take the following form

$$H = -\frac{1}{2} \sum_{1 \leq i \neq j \leq N} (q_i - q_j)^{-2} M_{ij}.\tag{23}$$

One can show that $[H, I_n] = 0$, $[I_n, I_m] = 0$, where $I_n = \sum_{j=1}^N (a_j^\dagger a_j)^n$, $a_j^\dagger = i \sum_{k=1}^N \frac{1}{q_j - q_k} M_{jk} + i q_j$, $a_j = (a_j^\dagger)^\dagger$, and $n, m = 1, 2, \dots, \infty$. These commutation relations yield integrability of the system. Having established the integrability, we have found the ground state of the model in the subspace of fixed number of holes, and fixed number of particles of each internal spin degree $N_\sigma, \sigma = 1, 2, \dots, \nu$ [13].

$$\phi(x_1\sigma_1, \dots, x_{N_e}\sigma_{N_e}; y_1, \dots, y_Q) = \prod_{i < j} (x_i - x_j)^{\delta_{\sigma_i\sigma_j}} e^{i\frac{\pi}{2} \text{sgn}(\sigma_i - \sigma_j)}. \quad (24)$$

It has been proved that this wavefunction has eigenenergy given by [13]

$$E_0 = -\frac{1}{4}N(N-1) + \frac{1}{2} \sum_{\sigma}^{\nu} \tilde{N}_{\sigma}(\tilde{N}_{\sigma} - 1). \quad (25)$$

Furthermore, we have found that the full energy spectrum of the system in the subspace of fixed number of particles of each internal spin degree can be written as [13]

$$E = E_0 + s \quad (26)$$

where $s = 0, 1, 2, \dots, s_{max}$, with an upper bound s_{max} due to the finite size of the Hilbert space. However, we have been unable to develop a systematic rule to characterize the degeneracy of the each energy level and to explain it with the underlying symmetries of the system.

VI. CONCLUSION AND ACKNOWLEDGEMENT

In this talk, we have reviewed some recent progresses in exact solutions of low dimensional electronic systems. We wish to thank Mo-lin Ge and F. Y. Wu for the invitation. This work was supported in part by the Swiss National Science Foundation.

REFERENCES

- [1] C. Gruber and N. Macris, *The Falicov-Kimball: review of rigorous results*, (1995). References therein.
- [2] E. H. Lieb and F. Y. Wu, Phys. Rev. Lett. **20**, 1445 (1968). C. N. Yang, Phys. Rev. Lett. **19**, 1312 (1967).
- [3] K. Michielson, Int. J. Mod. Phys. **B 7**, 2571 (1993).
- [4] E. H. Lieb, *The Hubbard model: some rigorous results and open problems*, in *Proceedings of the Conference “Advances in Dynamical Systems and Quantum Physics”* (World Scientific, Singapore, 1995).
- [5] F. D. M. Haldane, Phys. Rev. Lett. **60**, 635 (1988).
- [6] S. Shastry, Phys. Rev. Lett. **60**, 639 (1988).
- [7] Y. Kuramoto and H. Yokoyama, Phys. Rev. Lett. **67**, 1338 (1991).
- [8] N. Kawakami, Phys. Rev. B **45**, 7525 (1992).
- [9] D. F. Wang, James T. Liu and P. Coleman, Phys. Rev. B **46**, 6639 (1992).
- [10] A. P. Polychronakos, Phys. Rev. Lett. **70**, 2329 (1993).
- [11] M. Fowler and J. Minahan, Phys. Rev. Lett. **70**, 2325 (1993).
- [12] D. F. Wang and C. Gruber, Phys. Rev. B **49**, 15712 (1994).
- [13] C. Gruber and D. F. Wang, Phys. Rev. B **50**, 3103 (1994).
- [14] C. Gruber and D. F. Wang, invited talk at King’s College, May 1994.
- [15] B. Sutherland, Phys. Rev. A **5**, 1372 (1972), Phys. Rev. A **4**, 2019 (1971).
- [16] D. F. Wang and C. Gruber, Phys. Rev. B **51**, 7476 (1995).